



U.S. FDA Approves IBRANCE® (palbociclib) for the Treatment of Men with HR+, HER2- Metastatic Breast Cancer

Thursday, April 04, 2019 - 10:57am

Approval of expanded indication based predominately on real-world data

Pfizer (NYSE:PFE) today announced that the U.S. Food and Drug Administration (FDA) approved a supplemental New Drug Application (sNDA) to expand the indications for IBRANCE® (palbociclib) in combination with an aromatase inhibitor or fulvestrant to include men with hormone receptor-positive (HR+), human epidermal growth factor receptor 2-negative (HER2-) advanced or metastatic breast cancer. The approval is based on data from electronic health records and postmarketing reports of the real-world use of IBRANCE in male patients sourced from three databases: IQVIA Insurance database, Flatiron Health Breast Cancer database and the Pfizer global safety database.

“With this approval, we are now able to offer IBRANCE to the underserved male breast cancer community and provide more patients with HR+, HER2- metastatic breast cancer the opportunity to access an innovative medicine,” said Chris Boshoff, M.D., Ph.D., Chief Development Officer, Oncology, Pfizer Global Product Development. “We appreciate that our partnership with the FDA has allowed us to take a significant step forward in the use of real-world data to bring medicines to patients who are most in need.”

IBRANCE is now approved for adult patients with HR+, HER2- advanced or metastatic breast cancer in combination with an aromatase inhibitor as initial endocrine based therapy in postmenopausal women or in men; or with fulvestrant in patients with disease progression following endocrine therapy. With today’s approval, IBRANCE is the first and only CDK 4/6 inhibitor indicated in combination with an aromatase inhibitor for the first-

line treatment of men with HR+, HER2- metastatic breast cancer in the U.S.

“Men with breast cancer have limited treatment options, making access to medicines such as IBRANCE critically important,” said Bret Miller, founder of the Male Breast Cancer Coalition. “We applaud the use of real-world data, a new approach to drug review, to make IBRANCE available to certain men with metastatic breast cancer and help address an unmet need for these patients.”

Real-world data is playing an increasingly important role in expanding the use of already approved innovative medicines.¹ Due to the rarity of breast cancer in males, fewer clinical trials are conducted that include men resulting in fewer approved treatment options. In the U.S. in 2019, it is estimated that there will be 2,670 new cases of invasive breast cancer and about 500 deaths from metastatic breast cancer in males.² The 21st Century Cures Act, enacted in 2016, was created to help accelerate medical product development, allowing new innovations and advances to become available to patients who need them faster and more efficiently.³ This law places additional focus on the use of real-world data to support regulatory decision-making.⁴

Detailed analysis of the use of IBRANCE in men with HR+, HER2- advanced or metastatic breast cancer will be presented at an upcoming medical meeting. Based on limited data from postmarketing reports and electronic health records, the safety profile for men treated with IBRANCE is consistent with the safety profile in women treated with IBRANCE.⁵

About IBRANCE® (palbociclib) 125 mg capsules

IBRANCE is an oral inhibitor of CDKs 4 and 6,⁶ which are key regulators of the cell cycle that trigger cellular progression.^{7,8} In the U.S., IBRANCE is indicated for the treatment of adult patients with hormone receptor-positive (HR+), human epidermal growth factor receptor 2-negative (HER2-) advanced or metastatic breast cancer in combination with an aromatase inhibitor as initial endocrine based therapy in postmenopausal women or in men; or with fulvestrant in patients with disease progression following endocrine therapy.

IBRANCE currently is approved in more than 90 countries and has been prescribed to more than 200,000 patients globally.

The full prescribing information for IBRANCE can be found [here](#).

IMPORTANT IBRANCE® (palbociclib) SAFETY INFORMATION FROM THE U.S. PRESCRIBING INFORMATION

Neutropenia was the most frequently reported adverse reaction in PALOMA-2 (80%) and PALOMA-3 (83%). In PALOMA-2, Grade 3 (56%) or 4 (10%) decreased neutrophil counts were reported in patients receiving IBRANCE plus letrozole. In PALOMA-3, Grade 3 (55%) or Grade 4 (11%) decreased neutrophil counts were reported in patients receiving IBRANCE plus fulvestrant. Febrile neutropenia has been reported in 1.8% of patients exposed to IBRANCE across PALOMA-2 and PALOMA-3. One death due to neutropenic sepsis was observed in PALOMA-3. Inform patients to promptly report any fever.

Monitor complete blood count prior to starting IBRANCE, at the beginning of each cycle, on Day 15 of first 2 cycles and as clinically indicated. Dose interruption, dose reduction, or delay in starting treatment cycles is recommended for patients who develop Grade 3 or 4 neutropenia.

Based on the mechanism of action, IBRANCE can cause fetal harm. Advise females of reproductive potential to use effective contraception during IBRANCE treatment and for at least 3 weeks after the last dose. IBRANCE may impair fertility in males and has the potential to cause genotoxicity. Advise male patients to consider sperm preservation before taking IBRANCE. Advise male patients with female partners of reproductive potential to use effective contraception during IBRANCE treatment and for 3 months after the last dose. Advise females to inform their healthcare provider of a known or suspected pregnancy. Advise women not to breastfeed during IBRANCE treatment and for 3 weeks after the last dose because of the potential for serious adverse reactions in nursing infants.

The most common adverse reactions ($\geq 10\%$) of any grade reported in PALOMA-2 for IBRANCE plus letrozole vs placebo plus letrozole were neutropenia (80% vs 6%), infections (60% vs 42%), leukopenia (39% vs 2%), fatigue (37% vs 28%), nausea (35% vs 26%), alopecia (33% vs 16%), stomatitis (30% vs 14%), diarrhea (26% vs 19%), anemia (24% vs 9%), rash (18% vs 12%), asthenia (17% vs 12%), thrombocytopenia (16% vs 1%), vomiting (16% vs 17%), decreased appetite (15% vs 9%), dry skin (12% vs 6%), pyrexia (12% vs 9%), and dysgeusia (10% vs 5%).

The most frequently reported Grade ≥ 3 adverse reactions ($\geq 5\%$) in PALOMA-2 for IBRANCE plus letrozole vs placebo plus letrozole were neutropenia (66% vs 2%), leukopenia (25% vs 0%), infections (7% vs 3%), and anemia (5% vs 2%).

Lab abnormalities of any grade occurring in PALOMA-2 for IBRANCE plus letrozole vs placebo plus letrozole were decreased WBC (97% vs 25%), decreased neutrophils (95% vs 20%), anemia (78% vs 42%), decreased platelets (63% vs 14%), increased aspartate

aminotransferase (52% vs 34%), and increased alanine aminotransferase (43% vs 30%).

The most common adverse reactions ($\geq 10\%$) of any grade reported in PALOMA-3 for IBRANCE plus fulvestrant vs placebo plus fulvestrant were neutropenia (83% vs 4%), leukopenia (53% vs 5%), infections (47% vs 31%), fatigue (41% vs 29%), nausea (34% vs 28%), anemia (30% vs 13%), stomatitis (28% vs 13%), diarrhea (24% vs 19%), thrombocytopenia (23% vs 0%), vomiting (19% vs 15%), alopecia (18% vs 6%), rash (17% vs 6%), decreased appetite (16% vs 8%), and pyrexia (13% vs 5%).

The most frequently reported Grade ≥ 3 adverse reactions ($\geq 5\%$) in PALOMA-3 for IBRANCE plus fulvestrant vs placebo plus fulvestrant were neutropenia (66% vs 1%) and leukopenia (31% vs 2%).

Lab abnormalities of any grade occurring in PALOMA-3 for IBRANCE plus fulvestrant vs placebo plus fulvestrant were decreased WBC (99% vs 26%), decreased neutrophils (96% vs 14%), anemia (78% vs 40%), decreased platelets (62% vs 10%), increased aspartate aminotransferase (43% vs 48%), and increased alanine aminotransferase (36% vs 34%).

Avoid concurrent use of strong CYP3A inhibitors. If patients must be administered a strong CYP3A inhibitor, reduce the IBRANCE dose to 75 mg. If the strong inhibitor is discontinued, increase the IBRANCE dose (after 3-5 half-lives of the inhibitor) to the dose used prior to the initiation of the strong CYP3A inhibitor. Grapefruit or grapefruit juice may increase plasma concentrations of IBRANCE and should be avoided. Avoid concomitant use of strong CYP3A inducers. The dose of sensitive CYP3A substrates with a narrow therapeutic index may need to be reduced as IBRANCE may increase their exposure.

For patients with severe hepatic impairment (Child-Pugh class C), the recommended dose of IBRANCE is 75 mg. The pharmacokinetics of IBRANCE have not been studied in patients requiring hemodialysis.

About Pfizer Oncology

At Pfizer Oncology, we are committed to advancing medicines wherever we believe we can make a meaningful difference on the lives of patients. Today, Pfizer Oncology has an industry-leading portfolio of 18 approved innovative cancer medicines and biosimilars across more than 20 indications, including breast, prostate, kidney, lung and hematology. Pfizer Oncology is striving to change the trajectory of cancer.

Working together for a healthier world®

At Pfizer, we apply science and our global resources to bring therapies to people that extend and significantly improve their lives. We strive to set the standard for quality, safety and value in the discovery, development and manufacture of health care products. Our global portfolio includes medicines and vaccines as well as many of the world's best-known consumer health care products. Every day, Pfizer colleagues work across developed and emerging markets to advance wellness, prevention, treatments and cures that challenge the most feared diseases of our time. Consistent with our responsibility as one of the world's premier innovative biopharmaceutical companies, we collaborate with health care providers, governments and local communities to support and expand access to reliable, affordable health care around the world. For more than 150 years, we have worked to make a difference for all who rely on us. We routinely post information that may be important to investors on our website at www.pfizer.com. In addition, to learn more, please visit us on www.pfizer.com and follow us on Twitter at @Pfizer and @Pfizer_News, LinkedIn, YouTube, and like us on Facebook at [Facebook.com/Pfizer](https://www.facebook.com/Pfizer).

DISCLOSURE NOTICE: The information contained in this release is as of April 4, 2019. Pfizer assumes no obligation to update forward-looking statements contained in this release as the result of new information or future events or developments.

This release contains forward-looking information about the expansion of the indication for IBRANCE (palbociclib) in combination with an aromatase inhibitor or fulvestrant to include men with hormone receptor-positive (HR+), human epidermal growth factor receptor 2-negative (HER2-) advanced or metastatic breast cancer and Pfizer Oncology, including their potential benefits, that involves substantial risks and uncertainties that could cause actual results to differ materially from those expressed or implied by such statements. Risks and uncertainties include, among other things, uncertainties regarding the commercial success of IBRANCE; the uncertainties inherent in research and development, including the ability to meet anticipated clinical endpoints, commencement and/or completion dates for our clinical trials, regulatory submission dates, regulatory approval dates and/or launch dates, as well as the possibility of unfavorable new clinical data and further analyses of existing clinical data; the risk that clinical trial data are subject to differing interpretations and assessments by regulatory authorities; whether regulatory authorities will be satisfied with the design of and results from our clinical studies; whether and when any drug applications may be filed in any other jurisdictions for IBRANCE, for any additional indications for IBRANCE or for any other oncology products; whether and when any such applications may be approved by regulatory authorities, which will depend on myriad factors, including making a determination as to whether the product's benefits outweigh its known risks and determination of the

product's efficacy, and, if approved, whether IBRANCE or any such other oncology products will be commercially successful; decisions by regulatory authorities impacting labeling, safety, manufacturing processes and/or other matters that could affect the availability or commercial potential of IBRANCE or any other oncology products; and competitive developments.

A further description of risks and uncertainties can be found in Pfizer's Annual Report on Form 10-K for the fiscal year ended December 31, 2018 and in its subsequent reports on Form 10-Q, including in the sections thereof captioned "Risk Factors" and "Forward-Looking Information and Factors That May Affect Future Results," as well as in its subsequent reports on Form 8-K, all of which are filed with the U.S. Securities and Exchange Commission and available at www.sec.gov and www.pfizer.com.

1 FDA: Real World Evidence.

<https://www.fda.gov/scienceresearch/specialtopics/realworldevidence/default.htm>.

Accessed March 5, 2019. 2 American Cancer Society. Key statistics for breast cancer in men. <https://www.cancer.org/cancer/breast-cancer-in-men/about/key-statistics.html>.

Accessed March 20, 2019. 3 FDA: 21st Century Cures Act.

<https://www.fda.gov/regulatoryinformation/lawsenforcedbyfda/significantamendmentstothe fdca>

Accessed March 5, 2019. 4 FDA: Real World Evidence.

<https://www.fda.gov/ScienceResearch/SpecialTopics/RealWorldEvidence/default.htm>.

Accessed March 5, 2019. 5 Pfizer global safety database. 6 IBRANCE® (palbociclib)

Prescribing Information. New York, NY: Pfizer Inc: 2019. 7 Weinberg, RA. pRb and Control of the Cell Cycle Clock. In: Weinberg RA, ed. The Biology of Cancer. 2nd ed. New York, NY: Garland Science; 2014:275-329. 8 Sotillo E, Grana X. Escape from Cellular Quiescence. In: Enders GH, ed. Cell Cycle Deregulation in Cancer. New York, NY: Humana Press; 2010:3-22.

Pfizer Media: Jessica Smith 212-733-6213 Jessica.M.Smith@pfizer.com Pfizer Investors: Ryan Crowe 212-733-8160 Ryan.Crowe@pfizer.com